

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

2

1. AD-A216 477		1b. RESTRICTIVE MARKINGS	
2.		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited	
4. PERFORMING ORGANIZATION REPORT NUMBER(S)		5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR. 89-1868	
6a. NAME OF PERFORMING ORGANIZATION West Virginia University	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION AFOSR	
6c. ADDRESS (City, State, and ZIP Code) Cun-Quan Zhang, Mathematics Dept. West Virginia University, Morgantown, WV 26506		7b. ADDRESS (City, State, and ZIP Code) Bldg. 410 Bolling AFB, DC 20332 6448	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Air Force Office of Scientific Research	8b. OFFICE SYMBOL (If applicable) NM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR-89-0068	
8c. ADDRESS (City, State, and ZIP Code) AFOSR/NM Building 410 Bolling AFB, DC 20332-6448		10. SOURCE OF FUNDING NUMBERS	WORK UNIT ACCESSION NO.
		PROGRAM ELEMENT NO. AFOSR89-0068	TASK NO. A8
11. TITLE (Include Security Classification) Long Path Connectivity of Regular Graphs			
12. PERSONAL AUTHOR(S) Cun-Quan Zhang and Yong-Jin Zhu			
13a. TYPE OF REPORT Final Technical	13b. TIME COVERED FROM 11/88 TO 10/89	14. DATE OF REPORT (Year, Month, Day) 89/10/15	15. PAGE COUNT 17
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	Long Path Connectivity of Regular Graphs	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Any pair of vertices in a 4-connected, non-bipartite, k-regular graph are joined by a Hamilton path or a path of length at least 3k-6.			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL D. N. GLASSMAN		22b. TELEPHONE (Include Area Code) (202) 767-5026	22c. OFFICE SYMBOL ND


DTIC
ELECTE
JAN 05 1990
S DCS D

UNCLASSIFIED

00 01 04 161

LONG PATH CONNECTIVITY OF REGULAR GRAPHS

Cun-Quan Zhang*
 Department of Mathematics
 West Virginia University
 Morgantown, WV 26506



Yong-Jin Zhu
 Institute of System Science
 Academia Sinica
 Beijing, China

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

ABSTRACT: Any pair of vertices in a 4-connected non-bipartite k -regular graph are joined by a Hamilton path or a path of length at least $3k-6$.

*This research was partially supported by AFOSR under grant 89-0068

The topics about Hamilton cycles, circumferences and Hamiltonian connectivities of regular graphs have been interesting many mathematicians in recent years ([2],[1],[4],[7],[3],[6]).

In this paper, we will investigate the length of a longest path joining any pair of vertices of regular graphs and establish the following theorem.

THEOREM 1

Let G be a 4-connected non-bipartite k -regular graph. Then any pair of distinct vertices of G are joined by a Hamilton path or a path of length at least $3k-6$.

In a sense, this theorem is a generalization of the following results.

- (i) (Bollobas and Hobbs [1]) Any 2-connected k -regular graph of order at most $\frac{9}{4}k$ contains a Hamilton cycle.
- (ii) (Jackson [4]) Any 2-connected k -regular graph of order at most $3k$ contains a Hamilton cycle.
- (iii) (Zhu, Liu and Yu [7]) Any 2-connected k -regular graph of order at most $3k+3$ contains a Hamilton cycle.
- (iv) (Fan [3]) The length of a longest cycle in a 3-connected k -regular graph of order n is at least $\min\{n, 3k\}$.
- (v) (Zhang and Zhu [6]) Any pair of vertices of a 3-connected non-bipartite k -regular graph of order at most $3k-4$ are joined by a Hamilton path.

The condition of 4-connectivity in the theorem cannot be reduced. A 3-connected k -regular graph of order $3k+3$ containing no path of length at least $2k+3$ joining a pair of vertices can be constructed as follows. Let $k=3h$. Let G_1, \dots, G_9 be nine disjoint copies of complete graph K_h and

v_1, v_2, v_3 be three distinct vertices. Join an edge between each pair of vertices in $G_{3i+1} G_{3i+2} G_{3i+3}$ for $i=0,1,2$, and join an edge between v_j and each vertex of G_{3i+j} for $i=0,1,2$ and $j=1,2,3$. The induced graph contains $9h+3$ vertices and is $3h$ -regular 3-connected, in which v_i and v_j are not joined by any path of length longer than $6h+2$ for $i,j \in \{1,2,3\}$. (See fig. 1).

Actually, we can establish a result stronger than Theorem 1.

THEOREM 2. Let G be a 4-connected graph and x,y be a pair of distinct vertices of G such that

- (i) $d(v)=k$ for any vertex $v \in V(G) \setminus \{x,y\}$,
- (ii) $d(x), d(y) \leq k$.

Then the length of a longest path joining x and y is at least

- (i) $\min\{|V(G)|-1, 3k-6\}$ if G is not a bipartite graph, or G is a bipartite graph and x, y belong to different parts of the bipartition of G ;
- (ii) $\min\{|V(G)|-2, 3k-6\}$ if G is a bipartite graph and x,y belong to the same part of the bipartition of G .

Let $G=(V,E)$ be a graph with vertex set V and edge set E . Let $P=u_0 \cdots u_p$ be a path of G . For $0 \leq i, j \leq p$, the segment $u_i \cdots u_j$ of P is denoted by $u_i P u_j$ if $i \leq j$ or $u_i \bar{P} u_j$ if $i \geq j$. The length of a path P is the number of edges in P and is denoted by $l(P)$. Let H be a subgraph of G . Let w, w' be two vertices of H . The length of a longest

path of H joining w, w' is denoted by $L_H(w, w')$. Let v be a vertex of G . The set of vertices of H adjacent to v is denoted by $N_H(v)$ and the number of vertices of $N_H(v)$ is denoted by $d_H(v)$. When $V(H) = V(G)$, we simply write $d(v)$ and $N(v)$ instead of $d_G(v)$ and $N_G(v)$. Let $P = u_0 \cdots u_p$ be a path of G and X be a subset of $V(P)$. Denote

$$X^{+1} = \{u_{i+1} = u_i \in X\}$$

$$\text{and } X^{-1} = \{u_{i-1} = u_i \in X\}.$$

Let $E(H, H')$ be the set of all ordered pairs of vertices (x, y) such that $(x, y) \in E(G)$ and $x \in V(H)$, $y \in V(H')$. And let $|E(H, H')| = e(H, H')$. Note that if $V(H) \cap V(H') \neq \emptyset$, each edge (x, y) in the induced subgraph $G(V(H) \cap V(H'))$ will counted twice in $e(H, H')$ since the ordered pairs (x, y) and (y, x) are considered different in $E(H, H')$. Thus $d(v) = e(v, G)$ for any vertex v of G and $\sum_{v \in V(H)} d(v) = e(H, G)$ for subgraph H of G .

PROOF OF THEOREM 2

The theorem will be proved by contradiction. Suppose that the length of a longest path $P = v_0 \cdots v_p$ joining $x = v_0$ and $y = v_p$ is less than $3k-6$ and $G \setminus V(P)$ is not empty.

PART ONE. In this part, we will show that $G \setminus V(P)$ is an independent set of G . The following lemmas will be applied in this part.

LEMMA 1.1. (Lemma 4, [3]) Let H be a 2-connected graph and $Q = u_0 \cdots u_q$ be a longest path of H . Then

$$L_H(x, y) \geq \min\{d(u_0), d(u_q)\}$$

for any pair of distinct vertices x and y in H .

Let C be a set and $\{A_1, \dots, A_\alpha\}$, $\{B_1, \dots, B_h\}$ be partitions of C such that $\alpha \geq 2$ and $|A_\mu \cap B_j| \leq 1$ for any $\mu \in \{1, \dots, \alpha\}$ and any $j \in \{1, \dots, h\}$. If

$$B_i \cap A_\mu \neq \emptyset, B_j \cap A_\theta \neq \emptyset \text{ and } B_{i+1} = \dots = B_{j-1} = \emptyset$$

for some $\mu, \theta \in \{1, \dots, \alpha\}$ and $\mu \neq \theta$, then $\{i, \dots, j\}$ is called a closed extendible interval of $\{B_1, \dots, B_h\}$.

LEMMA 1.2 (Lemma 3.2, [6]) Let C be a set, $\{A_1, \dots, A_\alpha\}$ and $\{B_1, \dots, B_h\}$

be partitions of C defined as above. If s is an integer such that $\alpha \geq s$ and $|A_\mu| \geq s$ for each $\mu \in \{1, \dots, \alpha\}$, then $\{B_1, \dots, B_h\}$ has at least $s-1$ closed extendible intervals.

Suppose that $G \setminus V(P)$ is not an independent set and let W_0 be a component of $G \setminus V(P)$ which contains at least two vertices. Let T_1, \dots, T_t be all end-blocks of W_0 . (An end-block of W_0 is a block of W_0 which contains at most one cut-vertex of W_0).

I. We claim that there exists a longest path $Q_i = x_1^i \dots x_{q_i}^i$ in each T_i

such that

(i) $d_{W_0}(x_1^i) \leq d_{W_0}(x_{q_i}^i)$ and x_1^i is not a cut-vertex of W_0 , and

(ii) $d_{W_0}(x_1^i)$ is as big as possible.

Let $R = y_1 \dots y_r$ be a longest path in T_i such that $d_{W_0}(y_1) \leq d_{W_0}(y_r)$.

(α). If y_1 is a cut-vertex of W_0 and $d_{T_i}(y_1) \geq 2$, then there is another longest path $y_u \bar{R} y_1 y_{u+1} R y_r$ or $y_r \bar{R} y_{u+1} y_1 R y_u$ satisfying (i) for any $y_{u+1} \in N_R(y_1) \setminus \{y_2\}$. Of all longest paths in T_i satisfying (i), let $Q_i = x_1^i \cdots x_{q_i}^i$ be the one with the largest $d_{W_0}(x_1^i)$. (β) if y_1 is a cut-vertex of W_0 and $d_{T_i}(y_1) = 1$, then $|T_i| = 2$ and $R = y_1 y_2$ since T_i is a block. Hence $d_{W_0}(y_2) = 1$ and $d_{W_0}(y_1) > 1$ because y_1 is a cut-vertex of W_0 . It contradicts the assumption that $d_{W_0}(y_1) \leq d_{W_0}(y_r)$.

II. Let $d = \max\{d_{W_0}(x_i^1) : i = 1, \dots, t\}$. Without loss of generality, let $d = d_{W_0}(x_1^1)$.

(i) When $d \geq 2$ and $N_{Q_1}^{-1}(x_1^1) \cap \{\text{cut-vertices of } W_0\} = \emptyset$, let $Z = N_{Q_1}^{-1}(x_1^1)$.

(ii) When $d \geq 2$ and x_c^1 is a vertex of $N_{Q_1}^{-1}(x_1^1) \cap \{\text{cut-vertices of } W_0\}$.

Let $Z = [N_{Q_1}^{-1}(x_1^1) \setminus \{x_c^1\}] \cup \{x_1^2\}$.

In both cases (i) and (ii), we have that $|Z| = |N_{Q_1}^{-1}(x_1^1)| = d_{W_0}(x_1^1) = d$, and

by Lemma 1.1,

$$\begin{aligned}
L_{W_0}(z, z') &= L_{T_1}(z, z') \\
&\geq \min\{d_{T_1}(x_1^1), d_{T_1}(x_{q_1}^1)\} \\
&= d_{T_1}(x_1^1) \\
&= d_{W_0}(x_1^1) \\
&= d
\end{aligned}$$

for each pair of distinct vertices, $z, z' \in Z \setminus V(T_1)$. If $z \in Z \cap V(T_1)$ and

$z' \in Z \setminus T_1$ we have that $z' = x_1^2$ and

$$\begin{aligned}
L_{W_0}(z, z') &\geq L_{W_0}(z, x_c^1) + L_{W_0}(x_c^1, x_1^2) \\
&\geq L_{T_1}(z, x_c^1) \\
&\geq d
\end{aligned}$$

By the choice of Q_1 and x_1^1 , it follows that

$$d = d_{W_0}(x_1^1) \geq d_{W_0}(z)$$

for each $z \in Z$.

(iii) When $d=1$, T_1 is a single edge (x_1^1, x_2^1) . Hence, x_1^1 is a degree one vertex of W_0 and x_2^1 is either a cut-vertex of W_0 if $W_0 \neq T_1$, or a degree one vertex of W_0 if $W_0 = T_1$. If $W_0 = T_1$, then let $Z = \{x_1^1, x_2^1\}$. If

$W_0 \setminus T_1 \neq \emptyset$, by the choice of x_1^1 , we must have that $d_{W_0}(x_1^2) \leq d_{W_0}(x_1^1)$ and

x_1^2 is a degree one vertex of W_0 . Then let

$$Z = \{x_1^1, x_1^2\}.$$

Thus in either case, $d_{W_0}(z) = 1$ for any $z \in Z$.

So we always have that

$$|Z| = \max\{d, 2\}, \quad \dots \quad (1)$$

$$L_{W_0}(z, z') \geq d, \quad \dots \quad (2)$$

$$d_{W_0}(z) \leq d \text{ and } d_P(z) \geq k-d \quad \dots \quad (3)$$

for each pair of distinct vertices z and z' of Z . And

$$|T_1| \geq d+1 \quad \dots \quad (4)$$

since $d = d_{W_0}(x_1^1) = d_{T_1}(x_1^1)$.

III. We claim that $1 \leq d \leq k-4$.

Suppose that $d \geq k-3$. Since G is 4-connected, there are four intermediately disjoint paths $P_\mu = v_{i_\mu} \dots x_\mu$ joining T_1 and P for

$\mu = 1, \dots, 4$ where $\{v_{i_1}, v_{i_2}, v_{i_3}, v_{i_4}\}$ are distinct vertices of P ,

$0 \leq i_1 < i_2 < i_3 < i_4 \leq p$, $\{x_1, x_2, x_3, x_4\}$ belong to T_1 and

$$|\{x_1, \dots, x_4\}| = \min\{|T_1|, 4\}.$$

Let R_μ be a path joining x_μ and $x_{\mu+1}$ in T_1 such that R_μ is of length at least d if $x_\mu \neq x_{\mu+1}$ (by Lemma 1.1), or $R_\mu = x_\mu$ if $x_\mu = x_{\mu+1}$. Then

$$\ell(v_{i_\mu} P v_{i_{\mu+1}}) \geq \ell(v_{i_\mu} P x_\mu R_\mu x_{\mu+1} P_{\mu+1} v_{i_{\mu+1}}) \geq d+2$$

$$\text{if } x_\mu \neq x_{\mu+1}, \text{ or } \ell(v_{i_\mu} P v_{i_{\mu+1}}) \geq \ell(v_{i_\mu} P x_\mu P_{\mu+1} v_{i_{\mu+1}}) \geq 2$$

if $x_\mu = x_{\mu+1}$ since P is a longest path joining v_0 and v_p .

If $|T_1| \geq 4$, then $\{x_1, x_2, x_3, x_4\}$ are a set distinct vertices and

$$\begin{aligned} \ell(P) &\geq \sum_{\mu=1}^3 \ell(v_{i_\mu} P v_{i_{\mu+1}}) \\ &\geq 3(d+2) \\ &\geq 3k-3 \end{aligned} \quad (\text{by } d \geq k-3).$$

It contradicts the assumption that $\ell(P) < 3k-6$. Therefore $|T_1| \leq 3$ and

some x_i and x_j of $\{x_1, x_2, x_3, x_4\}$ are the same vertex. However,

$$\begin{aligned} 3k-7 &\geq \ell(P) \geq \sum_{\mu=1}^3 \ell(v_{i_\mu} P v_{i_{\mu+1}}) \\ &\geq \sum_{x_\mu \neq x_{\mu+1}} \ell(v_{i_\mu} P v_{i_{\mu+1}}) + \sum_{x_\mu = x_{\mu+1}} \ell(v_{i_\mu} P v_{i_{\mu+1}}) \\ &\geq (d+2)(|T_1| - 1) + 2(4 - |T_1|) \\ &= d(|T_1| - 1) + 6 \\ &\geq d^2 + 6 \quad (\text{by (4)}) \\ &\geq k^2 - 6k + 15 \quad (\text{by } d \geq k-3). \end{aligned}$$

Thus $0 \geq k^2 - 9k + 22$. But the value of $k^2 - 9k + 22$ is always positive for any k .

It leads a contradiction and follows our claim.

IV. Now we wish to show the following inequality

$$l(P) \geq (k-d-1)(d+2) \quad \dots \dots \dots (5)$$

Let z, z' be a pair of distinct vertices of Z . We have known that

$d_P(z), d_P(z') \geq k - d$ and $L_{W_0}(z, z') \geq d$ (by (2) and (3)). Let

$|N_P(z) \cap N_P(z')| = \sigma(z, z')$. Since P is a longest path joining v_0 and

v_p , $N_P(z) \cup N_P(z')$ does not contain two consecutive vertices of P . Let

$\{v_{i_1}, \dots, v_{i_r}\} = N_P(z) \cup N_P(z')$. Then $[v_{i_1} P v_{i_r}] \setminus [N_P(z) \cup N_P(z')]$ contains $r-1$

open segments. A segment $v_{i_\theta} P v_{i_{\theta+1}}$ is called extendible with respect to

$\{z, z'\}$ if either $v_{i_\theta} \in N(z)$ and $v_{i_{\theta+1}} \in N(z')$ or $v_{i_\theta} \in N(z')$ and $v_{i_{\theta+1}} \in N(z)$.

Otherwise, it is called unextendible. It is not very hard to see that P has at least $\sigma(z, z') - 1$ extendible segments with respect to $\{z, z'\}$. Since P is a longest path joining v_0 and v_p and $L_{W_0}(z, z') \geq d$, each extendible segment is of length at least $d+2$ and each unextendible segment is of length at least two.

(i) If there is a pair of distinct vertices $\{z_1, z_2\}$ of Z such that P has $\sigma(z_1, z_2)$ or $\sigma(z, z) - 1$ extendible segments with respect to

$\{z_1, z_2\}$ then one of $\{N_P(z_1), N_P(z_2)\}$ must be a subset of another one and

$$\sigma(z_1, z_2) = \min(|N_P(z_1)|, |N_P(z_2)|) \geq k-d.$$

$$\begin{aligned} \text{So } l(P) &\geq (\text{total length of all extendible segments}) \\ &\geq (d+2)(\sigma(z_1, z_2)-1) \\ &\geq (d+2)(k-d-1). \quad (\text{since } \sigma(z_1, z_2) \geq k-d) \end{aligned}$$

Thus we have established the inequality (5) in this case, and therefore we will assume that P has at least $\alpha(z, z') + 1$ extendible segments with respect to any pair of distinct vertices $\{z, z'\}$ of Z .

$$(ii) \text{ Case 1. } d \leq \frac{k}{2}$$

Let $\sigma = \max \{\sigma(z, z') \mid z, z' \text{ are a pair of distinct vertices of } Z\}$.

Choose a pair of distinct vertices z_1 and z_2 of Z such that $\sigma(z_1, z_2) = \sigma$

and let $r = |N_P(z_1) \cup N_P(z_2)|$. It is clear that

$$r + \sigma = |N_P(z_1)| + |N_P(z_2)| \geq 2(k-d) \quad \dots\dots\dots (6)$$

$$r \geq |N_P(z_1)| \geq k-d \quad \dots\dots\dots (7)$$

Since P has at least $\sigma+1$ extendible segments with respect to $\{z_1, z_2\}$, we

have that

$$\begin{aligned} l(P) &\geq (\text{total length of all extendible segments with} \\ &\quad \text{respect to } \{z_1, z_2\} + \\ &\quad (\text{total length of all unextendible segments with} \\ &\quad \text{respect to } \{z_1, z_2\})) \\ &\geq (d+2)(\sigma+1) + 2[(r-1) - (\sigma+1)] \\ &= 2r + \sigma d + d - 2 \\ &\geq 2[2(k-d) - \sigma] + \sigma d + d - 2 \quad (\text{since } r \geq 2(k-d) - \sigma \text{ by (6)}) \end{aligned}$$

$$= 4k - 4d - 2\sigma + \sigma d + d - 2$$

$$= (4k - 2d) - 2d + (\sigma + 1)(d - 2)$$

$$\geq 3k - 2d + (\sigma + 1)(d - 2) \quad (\text{since } d \leq \frac{k}{2})$$

$$\text{Thus } 3k - 7 \geq \ell(P) \geq 3k - 2d + (\sigma + 1)(d - 2) \quad \dots\dots\dots (8)$$

if $\sigma \geq 1$, by (8), we have that

$$\begin{aligned} 3k - 7 &\geq 3k - 2d + 2(d - 2) \\ &= 3k - 4. \end{aligned}$$

It is a contradiction and hence we have that $\sigma = 0$. If $d \leq 4$, by (8), we have that

$$\begin{aligned} 3k - 7 &\geq \ell(P) \geq 3k - 2d + (d - 2) && (\text{since } \sigma = 0) \\ &\geq 3k - 6 && (\text{since } d \leq 4). \end{aligned}$$

It is also a contradiction and therefore we must have that $d \geq 5$. Note that $|Z| \geq d \geq 5$, let z, z', z'' be three distinct vertices of Z . By the

definition of σ and $\sigma = 0$, the subsets $N_P(z)$, $N_P(z')$ and $N_P(z'')$ of $V(P)$ are pairwise disjoint. Hence

$$|N_P(z) \cup N_P(z') \cup N_P(z'')| \geq 3(k - d)$$

and P has at least $3(k - d) - 1$ segments each of which is of length at least two. So

$$\begin{aligned} \ell(P) &\geq 2[3(k - d) - 1] \\ &= 6k - 6d - 2 \\ &\geq 3k - 2 && (\text{since } d \leq \frac{k}{2}). \end{aligned}$$

It contradicts that $\ell(P) \leq 3k - 7$.

(iii) Case 2. $d \geq \frac{k}{2}$.

Let $C = E(Z, P)$

be a set and

$$\{A_z = E(z, P) : \text{for each } z \in Z\}$$

and $\{B_i = E(Z, v_i) : \text{for each } v_i \in V(P)\}$

be partitions of C . Note that $|\{A_z\}| = |Z| = d \geq k-d$ and $|A_z| = d_P(z) \geq k-d$

for any $z \in Z$ (by (3)), $|A_z \cap B_i| \leq 1$ for any $z \in Z$ and $v_i \in V(P)$. We can apply

Lemma 1.2 on C and these two partitions of C . Thus P has at least $k-d-1$ extendible segments each of which is of length at least $d+2$ and therefore

$$\begin{aligned} \ell(P) &\geq (\text{total length of all extendible segments}) \\ &\geq (d+2)(k-d-1) \end{aligned}$$

and the inequality (5) holds for all cases.

V. Since $1 \leq d \leq k-4$, the minimum value of $(d+2)(k-d-1)$ is $3k-6$ it contradicts that $\ell(P) < 3k-6$ and therefore, $G \setminus V(P)$ is an independent set.

Part two.

It has been shown in part one that $W = G \setminus V(P)$ is an independent set.

Let $w \in W$. Following [5], put $Y_0 = \phi$ and for $i \geq 1$, put

$$X_i = N(Y_{i-1} \cup \{w\})$$

and $Y_i = \{v_j \in V(P) : v_{j-1} \in X_i \text{ and } v_{j+1} \in X_i\}$.

Thus $N(w) = X_1 \subseteq X_2 \subseteq \dots$ and $\phi = Y_0 \subseteq Y_1 \subseteq Y_2 \subseteq \dots$.

Put $X = \bigcup_{i=1}^{\infty} X_i$ and $Y = \bigcup_{i=1}^{\infty} Y_i$. The follow lemma has been proved in [6] and

will be applied in this part of the proof.

LEMMA 2.1.

(i) (direct conclusion of the definition) $Y \subseteq V(P) \setminus \{v_0, v_p\}$ and

$$Y = (X \cap P)^{+1} \cup (X \cap P)^{-1}.$$

(ii) (Lemma 4.4. [6]) X does not contain two consecutive vertices of P .

(iii) (Lemma 4.4. [6]) $X \cap Y = \emptyset$.

(iv) (Lemma 4.7. [6]) $Y \cup W$ is an independent set of G , $N(Y) \subseteq V(P)$ and $N(Y \cup \{w\}) = X \subseteq V(P)$.

(v) $e(X, Y \cup \{w\}) = k(|Y| + 1)$ and $e(V', Y \cup \{w\}) = 0$ for any subset V' of $V(G) \setminus X$.

Proof. We only need to prove (v). By (i) $v_0, v_p \notin Y \cup \{w\}$, it follows that

$d(u) = k$ for any $u \in Y \cup \{w\}$. Since $X = N(Y \cup \{w\})$, $e(Y \cup \{w\}, X) = e(Y \cup \{w\}, G) = k|Y \cup \{w\}|$ and $N(Y \cup \{w\}) \cap V' = \emptyset$ for any subset V' of $V(G) \setminus X$.

Put $|X| = \chi$ and $|Y| = \psi$. Then $P \setminus X \cup Y$ is a union of at most $\chi + \psi + 1$ segments of P . Let S_1, \dots, S_{t-1} be the segments of $P \setminus X \cup Y$ not containing v_0 and v_p . Let S_0 (or S_t) be the segment of $P \setminus X \cup Y$ containing v_0 (or v_p , respectively) if v_0 (or v_p , respectively) does not belong to X .

Obviously, $S_0 = \emptyset$ (or $S_t = \emptyset$) if $v_0 \in X$ (or $v_p \in X$, respectively). It is easy

to see that $|S_i| \geq 2$ for $1 \leq i \leq t-1$ and $t = \chi - \psi$. Let $S = \bigcup_{i=0}^t S_i$. Here

Here $V(P) = XuYuS$, by (i) and (iv) of Lemma 2.1.

Case 1. $S \neq \emptyset$.

Let $Z_i = S_i \cap (X^{+1}uX^{-1})$ and $Z = \bigcup_{i=0}^t Z_i$. We have that

LEMMA 2.2 (Lemma 4.8, [6])

$$e(Z, S) \leq (t - \lambda)(|S| - t + 3)$$

where $\lambda = 0$ if $S_0 u S_t \neq \emptyset$ and $\lambda = 1$ if $S_0 u S_t = \emptyset$.

and

LEMMA 2.3 (Lemma 4.9, [6])

$$e(X, W \setminus \{w\}) \geq e(Z, W \setminus \{w\}).$$

Now we can prove our theorem in this case. Since

$$k\chi \geq e(X, G) \geq e(X, Z) + e(X, Yu\{w\}) = e(X, W \setminus \{w\})$$

and

$$k|Z| = e(Z, G) = e(Z, X) + e(Z, Yu\{w\}) + e(Z, S) + e(Z, W \setminus \{w\}),$$

we have that

$$\begin{aligned} k\chi - e(X, Yu\{w\}) - e(X, W \setminus \{w\}) &\geq e(X, Z) \\ &= e(Z, X) \\ &= k|Z| - e(Z, S) - e(Z, W \setminus \{w\}) - e(Z, Yu\{w\}). \end{aligned}$$

Thus

$$\begin{aligned} k\chi - k(\psi + 1) - e(X, W \setminus \{w\}) \\ \geq k|Z| - e(Z, S) - e(Z, W \setminus \{w\}) \end{aligned}$$

by (v) of Lemma 2.1. Note that $\chi - \psi = t$ and

$$e(X, W \setminus \{w\}) \geq e(Z, W \setminus \{w\})$$

(by Lemma 2.3), it follows that

$$e(Z, S) \geq -kt + k + k |Z|.$$

When $S_0 \cup S_t \neq \emptyset$, $|Z| \geq 2t-1$. By Lemma 2.2,

$$t(|S| - t + 3) \geq -kt + k + k(2t-1).$$

Simplifying the above inequality, we have that

$$|S| \geq t-3+k. \quad \dots \quad (9)$$

When $S_0 \cup S_t = \emptyset$, $|Z| = 2(t-1)$. By Lemma 2.2,

$$(t-1)(|S| - t + 3) \geq -kt + k + 2k(t-1).$$

Simplifying the above inequality, we obtain the inequality (9) again. Since

$V(P) = S \cup X \cup Y$, and $t+4 = \chi \geq |N(w)| = k$,

$$\begin{aligned} l(P)+1 &= |V(P)| = |S| + |X| + |Y| \\ &\geq (t-3+k) + \chi + \psi \quad (\text{by (9)}) \\ &= k + 2\chi - 3 \\ &\geq 3k - 3 \end{aligned}$$

It contradicts that $l(P) < 3k-6$ and therefore the path joining v_0 and v_p is of length at least $3k-6$ in the case of $S \neq \emptyset$.

Case two. $S = \emptyset$. In this case, we must have $p = l(P)$ is even and

$X = \{v_{2i} : i=0, \dots, \frac{p}{2}\}$, $Y = \{v_{2i-1} : i=1, \dots, \frac{p}{2}\}$. Thus $|Y \cup \{w\}| = |X|$. We claim

that X is also an independent set and $N(X) \subseteq Y \cup \{w\}$. By (v) of Lemma 2.1, we have that

$$e(Y \cup \{w\}, X) = k |Y \cup \{w\}| = k |X|.$$

Since the maximum degree of G is k , all neighbors of every vertex of X are contained in $Y \cup \{w\}$.

Moreover, by (iv) of Lemma 2.1, both X and $Y \cup \{w\}$ are independent sets and

$$E(X, Y \cup \{w\}) = E(X, G) = E(G, Y \cup \{w\}).$$

The connectivity of G implies that $V(G) = XuYu\{w\}$. Thus $(X, Y \cup \{w\})$ is a bipartition of G and v_0, v_p are joined by a path of length $|V(G)| - 2$.

REFERENCES

- [1]. Bollobas, B. and Hobbs, A. M., Hamiltonian cycles in regular graphs, in "Advances in Graph Theory", North Holland, Amsterdam, 1978.
- [2]. Erdos, P. and Hobbs, A. M., Hamiltonian cycles in regular graphs of moderate degree, J. Combinatorial Theory, B. 23(1977) p.139-142.
- [3]. Fan, G., Longest cycle in regular graphs, J. Combinatorial Theory, B.39 (1985) p.325-345.
- [4]. Jackson, B., Hamilton cycles in regular 2-connected graphs, J. Combinatorial Theory, B. 29 (1980), p.27-46.
- [5]. Woodall, D. R., The binding number of a graph and its Anderson number, J. Combinatorial Theory, B. 15 (1973), p.225-255.
- [6]. Zhang, C. Q. and Zhu, Y. J., Hamiltonian connectivity and factorization.
- [7]. Zhu, Y. J., Liu, Z. H., and Yu, Z. G., 2-connected k -regular graphs on at most $3k+3$ vertices to be Hamiltonian, J. System Sciences & Math. Sciences, 6(1986), p.36-49, p.136-145.




fig. 1.